arXiv:cond-mat/0410441v1 [cond-mat.stat-mech] 18 Oct 2004

Thermal-equilibrium properties of one-dimensional gas with short-ranged repulsion

Milan Krbalek
Faculty of Nuclear Sciences and Physical Engineering,
Czech Technical University,
Trojanova 13, 120 00 Prague, Czech Republic*

February 2, 2008

Abstract

We derive the exact formula for thermal-equilibrium spacing distribution of one-dimensional particle gas with repulsive potential depending on the distance r between the neighboring particles. We are focused on the power-law potentials $r^{-\alpha}$ for $\alpha > 0$, being motivated by the actual traffic research.

PACS numbers: 05.70-a, 05.20-y

Investigation of one-dimensional particle ensembles seems actually to be very useful for understanding of the complex system called *vehicular traffic.* Beside the popular cellular automata, in the recent time new trend appears in the traffic modelling. Application of the equilibrium statistical-physics to non-equilibrium traffic systems has been successfully demonstrated for example in the Ref.[1]. In this continuing work we analyze a certain family of the particle gases exposed to the heat bath with the temperature $T \geq 0$, being primarily aimed at the equilibrium distribution for particle distance (*spacing distribution*) and particle velocity (*velocity distribution*). Subsequently, such a detailed description can be used for comparing with the microscopic structure of the realistic traffic flows (in Ref.[1]).

Consider N identical particles on the circle of the circumference $L = N$. Let x_i $(i = 1, 2, \ldots, N)$ represents the circular position of the $i-$th particle. Put $x_{N+1} = x_1 + 2\pi$, for convenience. Now we introduce the short-ranged potential energy

$$U \propto \sum_{i=1}^{N} V\left(r_i\right),$$

where $V(r_i)$ corresponds to the repulsive two-body potential depending on the distance $r_i = |x_{i+1} - x_i|\frac{N}{2\pi}$ between the neighboring particles only. Thus, the hamiltonian of the described ensemble reads as

$$\mathcal{H} = \frac{1}{2}\sum_{i=1}^{N}(v_i - \overline{v})^2 + C\sum_{i=1}^{N} V\left(r_i\right),$$

with the $i-$th particle velocity v_i and the positive constant C. Note that $\overline{v}$ represents the mean velocity in the ensemble. Then, the appropriate partition function[1]

$$\mathcal{Z} = \int_{\mathbb{R}^{2N}} \delta\left(L - \sum_{i=1}^{N} r_i\right) e^{-\frac{\mathcal{H}}{kT}}\; dr_1 dv_1 \ldots dr_N dv_N \tag{1}$$

leads us to the simple assertion that velocity v of particles is Gaussian distributed, i.e.

$$P(v) = \frac{1}{\sqrt{2\pi kT}}\; e^{\frac{(v-\overline{v})^2}{2kT}}.$$

is the corresponding probability density.

Of larger interest, however, is the spacing distribution $P_\beta(r)$. In order to calculate the exact form of $P_\beta(r)$, one can restrict the partition function (1) to the reduced form

$$\mathcal{Z}_N(L) = \int_{\mathbb{R}^N} \delta\left(L - \sum_{i=1}^{N} r_i\right) e^{-\beta\sum_{i=1}^{N} V(r_i)}\, dr_1 \ldots dr_N$$

---

*Electronic address: `milan.krbalek@fjfi.cvut.cz`

[1] k represents the Boltzmann factor

where $\beta = \frac{C}{kT}$ is so-called *inverse temperature* of the heat bath. Denoting $f(r) = e^{-\beta V(r)}$ the previous expression changes to

$$\mathcal{Z}_N(L) = \int_{\mathbb{R}^N} \delta\left(L - \sum_{i=1}^{N} r_i\right) \prod_{i=1}^{N} f(r_i)\ dr_1 \ldots dr_N.$$

Applying the Laplace transformation (see the Ref.[2] for details) one can obtain

$$g_N(p) \equiv \int_0^\infty \mathcal{Z}_N(L)\ e^{-pL}\, dL =$$

$$= \left(\int_0^\infty f(r) e^{-pr}\, dr\right)^N \equiv [g(p)]^N.$$

Then the partition function (in the large N limit) can be computed with the help of Laplace inversion

$$\mathcal{Z}_N(L) = \frac{1}{2\pi \mathtt{i}} \int_{B-\mathtt{i}\infty}^{B+\mathtt{i}\infty} g_N(p)\ e^{Lp}\ dp.$$

Its value is well estimated by the approximation in the saddle point B which is determined using the equation

$$\frac{1}{g(B)} \frac{\partial g}{\partial p}(B) = -\frac{L}{N}.$$

Thus,

$$\mathcal{Z}_N(L) \approx [g(B)]^N\, e^{LB}. \tag{2}$$

Hence the probability density for spacing r_1 between the particles ♯2 and ♯1 can be then reduced to the form

$$P(r_1) = \frac{\mathcal{Z}_{N-1}(L - r_1)}{\mathcal{Z}_N(L)} f(r_1).$$

Supposing $N \gg 1$ and using equation (2) we obtain

$$P(r_1) = \frac{1}{g(c)} f(r_1)\ e^{-Br_1},$$

which leads to the distribution function for spacing r between arbitrary couple of neighboring particles

$$P_\beta(r) = A\ e^{-\beta V(r)}\ e^{-Br}. \tag{3}$$

Note that constant A assures the normalization $\int_0^\infty P_\beta(r)\ dr = 1$. Furthermore, returning to the original choice $L = N$, the mean spacing is

$$\langle r \rangle \equiv \int_0^\infty r P_\beta(r)\ dr = 1. \tag{4}$$

Two above conditions can be understood as equation system for unknown normalization constants A, B.

Let us to proceed to the special variants of the gas studied. Firstly, we draw our attention to the Coulomb gas with the logarithmic potential

$$V(r) := -\ln(r).$$

Such a gas is frequently used in the many branches of physics (including the traffic research in Ref. [4]) and the corresponding spacing distribution reads as (see Ref. [2])

$$P_\beta(r) = \frac{(\beta+1)^{\beta+1}}{\Gamma(\beta+1)}\ r^\beta e^{-(\beta+1)r},$$

where $\Gamma(\xi)$ is gamma function. Of larger physical interest, as demonstrated in Ref. [1] and [3], seems actually to be the potential

$$V_\alpha(r) := r^{-\alpha},$$

for $\alpha > 0$. The aim of the following computational procedure is to normalize the distribution

$$P_\beta(r) = A e^{-\frac{\beta}{r^\alpha}} e^{-Br}. \tag{5}$$

Consider now the favorable choice $\alpha = 1$, for which the normalization integrals are exactly expressed as

$$\begin{aligned} \int_0^\infty e^{-\frac{\beta}{r}} e^{-Br}\ dr &= 2\sqrt{\frac{\beta}{B}}\ \mathcal{K}_1\left(2\sqrt{\beta B}\right) \\ \int_0^\infty r e^{-\frac{\beta}{r}} e^{-Br}\ dr &= 2\frac{\beta}{B}\ \mathcal{K}_2\left(2\sqrt{\beta B}\right), \end{aligned} \tag{6}$$

where $\mathcal{K}_\lambda$ is the Mac-Donald's function (modified Bessel's function of the second kind) of order λ, having for $\lambda = 1$ and $\lambda = 2$ an approximate expression

$$\mathcal{K}_\lambda(y) = \sqrt{\frac{\pi}{2}} e^{-y} \left(y^{-1/2} + \frac{3}{8} 5^{\lambda-1} y^{-3/2} + \mathcal{O}\left(y^{-5/2}\right)\right).$$

Applying the equations (6) to the normalization integrals one can determine the exact values of the constants A and B. Both of them can be, moreover, very well estimated by the large β approximations

$$B \approx \frac{900\ \beta}{\left(3 - 16\beta + \sqrt{9 + 864\beta + 256\beta^2}\right)^2} \approx \beta + \frac{3}{2}$$

and

$$A \approx \frac{1}{2}\sqrt{1 + \frac{3}{2\beta}} \frac{1}{\mathcal{K}_1\left(\sqrt{2\beta(2\beta+3)}\right)}.$$

Finally, we investigate the distribution (5) for general $\alpha > 0$. Although in this case the normalization integrals are not trivially solvable, the scaling (4) leads us to the simple formula

$$B \approx \alpha\beta + 1 + \frac{\alpha}{2} \qquad (\beta \to \infty)\,. \tag{7}$$

The rough estimation $r^{-\alpha} \approx 1-\alpha+\alpha r^{-1}$ provides the asymptotical formula for normalization constant A :

$$A \approx \frac{1}{2}\sqrt{1+\frac{1}{2\beta}+\frac{1}{\alpha\beta}}\,\frac{e^{\beta(1-\alpha)}}{\mathcal{K}_1\left(\sqrt{2\alpha\beta(2\alpha\beta+\alpha+2)}\right)}. \tag{8}$$



Figure 1. Relative deviation in the approximate value of the normalization constant B as function of parameter β. We display the deviation (10) between the numerical value B_{ex} and the value B_{est}, obtained from the large β approximation (7). The plus signs, squares, triangles, circles and stars correspond to the parameters $\alpha = 1, 2, 3, 4, 5$, respectively. The tails of the curves are magnified on the insert.

For practical applications it seems to be useful to detect the critical inverse temperature β_{crit} under which the relative deviation between the exact (ex) and estimated (est) values of the constant A (or B)

$$\delta_{\log(A)} := \frac{|\log(A_{ex)} - \log(A_{est})|}{\log(A_{ex})} \tag{9}$$

$$\delta_B := \frac{|B_{ex} - B_{est}|}{B_{ex}} \tag{10}$$

are larger then the fixed permissible deviation δ. For these purposes we plot the functional dependence $\delta_B = \delta_B(\beta)$ and $\delta_{\log(A)} = \delta_{\log(A)}(\beta)$ in the Figure 1 and Figure 2, respectively. We note that the exact values A_{ex}, B_{ex} were determined with the help of the numerical computations.



Figure 2. Relative deviation in the approximate value of the normalization constant A as function of parameter β. Plotted is the deviation (9) between the numerically computed value A_{ex} and the estimated value (8). The symbols used here are consistent with the symbols in the Fig. 1.

To conclude, we have found the exact form of the thermal-equilibrium spacing distribution for one-dimensional gas which neighboring particles are repulsed by the two-body potential $V = r^{-\alpha}$, where r is their mutual distance. The values of two normalization constants were successfully estimated by the simple approximation. The determined distribution has been and will be applied in the traffic research where compared to the distance clearance distribution of the freeway traffic samples.

References

[1] M. Krbalek and D. Helbing: *Determination of interaction potentials in freeway traffic from steady-state statistics*, Physica A **333** (2004), 370 – 378

[2] E.B. Bogomolny, U. Gerland, and C. Schmit: *Short-range plasma model for intermediate spectral statistics*, Eur. Phys. J. B **19** (2001), 121 – 132

[3] P. Seba and M. Krbalek, *Statistical properties of the city transport in Cuernavaca (Mexico) and random matrix ensembles,* J. Phys. A **33** (2000), L229 – L234

[4] M. Krbalek, P. Seba, and P. Wagner, *Headways in traffic flow: Remarks from a physical pescpective,* Phys. Rev. E **64** (2001), 066119 – 066125